


Barren River Area Development District (ADD) 2017 Arthritis Fact Sheet

Arthritis Prevalence by ADD, 2015 KyBRFS



How many Kentuckians in the Barren River ADD have Arthritis?

- Over 73,000 adults (32%) ages 18 and over¹
- Women (35%); Men (29%)¹
- Higher among those earning less than \$25,000 annually (48%)¹

How does Arthritis impact daily life in Barren River ADD?

- 39,151 reported limits to daily activities¹
- 32,980 working-age adults (18-64) reported work limitations¹
- 18,525 adults restricted social activities due to symptoms¹

How many in the Barren River ADD have had knee or hip replacement?

- 871 adults hospitalized for knee arthroplasty²
 - ⇒ average length of stay was 3.1 days²
 - ⇒ average charges per stay were \$14,862²
- 523 adults hospitalized for hip replacement (total or partial)²
 - ⇒ average length of stay was 3.6 days²
 - ⇒ average charges per stay were \$14,640²

1. Kentucky Behavioral Risk Factor Survey (KyBRFS) 2015 Data. Kentucky Department for Public Health, Cabinet for Health and Family Services, Frankfort, Kentucky

2. HCUPnet, Healthcare Cost and Utilization Project. Agency for Healthcare Research and Quality, Rockville, MD. <https://hcupnet.ahrq.gov/>. Accessed January 5, 2017

What can be done to improve Arthritis in Barren River ADD?

Individuals

Low impact physical activity can lessen arthritis pain and stiffness. Proven self-management and physical activity programs such as the Chronic Disease Self-Management Program, Enhance Fitness, and Walk with Ease are available in most areas of Kentucky through your Area Agency on Aging and Independent Living, local health department, Senior Center, or YMCA.

Health Care Providers

Prescribe low impact physical activity appropriate for those with arthritis. Refer patients to evidence-based interventions such as those listed above to help reduce symptoms and improve health and overall quality of life.

